

SHIP CQM Measures #5 - 7

Measure Title		Current SHIP Language	HIT Workgroup Recommended Changes		Comments	
Childhood Immunization Status	Percentage of children two years of age who had four DtaP/DT, three IPV, one MMR, three H influenza type B, three hepatitis B, one chicken pox vaccine, and four pneumococcal conjugate vaccines by their second birthday. The measure calculates a rate for each vaccine and two separate combination rates.	Percentage of children 2 years of age who had four diphtheria, tetanus and acellular pertussis (DTaP); three polio (IPV), one measles, mumps and rubella (MMR); three H influenza type B (HiB); three hepatitis B (Hep B); one chicken pox (VZV); four pneumococcal conjugate (PCV); one hepatitis A (Hep A); two or three rotavirus (RV); and two influenza (flu) vaccines by their second birthday CMS117v5 (2017), PQRS 240, NQF 0038	Numerator	Children who have evidence showing they received recommended vaccines, had documented history of the illness, had a seropositive test result, or had an allergic reaction to the vaccine by their second birthday	It appears that the SHIP measure was previously aligned with the national measure. The recommended changes reflect the updates to the national measure.	
			Denominator	Children who turn 2 years of age during the measurement period and who have a visit during the measurement period		
Preventive Care and Screening: Screening for Depression and Follow-Up Plan	Percentage of patients aged 12 years and older screened for clinical depression using a standardized tool and follow up plan documented.	Percentage of patients aged 12 years and older screened for depression on the date of the encounter using an age appropriate standardized depression screening tool AND if positive, a follow-up plan is documented on the date of the positive screen CMS2v6 (2017), PQRS 134, NQF 0418	Numerator	Patients screened for depression on the date of the encounter using an age appropriate standardized tool AND if positive, a follow-up plan is documented on the date of the positive screen	It appears that the SHIP measure was previously aligned with the national measure. The recommended changes reflect the updates to the national measure.	
			Denominator	All patients aged 12 years and older before the beginning of the measurement period with at least one eligible encounter during the measurement period		
Non-Malignant Opioid Use	Percent of patients chronically prescribed an opioid medication for non-cancer pain (defined as three consecutive months of prescriptions) that have a controlled substance agreement in force (updated annually).	Opioid Therapy Follow-up Evaluation - All patients 18 and older prescribed opiates for longer than six weeks duration who had a follow-up evaluation conducted at least every three months during Opioid Therapy documented in the medical record. (PQRS 408)	Numerator	Patients who had a follow-up evaluation conducted at least every three months during opioid therapy	The Opioid Rx will likely be in the EMR meds list; however, it may be hard to know if they have been taking them for 6 weeks	
			Denominator	All patients 18 and older prescribed opiates for longer than six weeks duration		
			Documentation of Signed Opioid Treatment Agreement - All patients 18 and older prescribed opiates for longer than six weeks duration who signed an opioid treatment agreement at least once during Opioid Therapy documented in the medical record. (PQRS 412)	Numerator	Patients who signed an opioid treatment agreement at least once during opioid therapy	The Opioid Rx will likely be in the EMR meds list; however, it may be hard to know if they have been taking them for 6 weeks
				Denominator	All patients 18 and older prescribed opiates for longer than six weeks duration	
			Evaluation or Interview for Risk of Opioid Misuse - All patients 18 and older prescribed opiates for longer than six weeks duration evaluated for risk of opioid misuse using a brief validated instrument (e.g. Opioid Risk Tool, SOAAP-R) or patient interview documented at least once during Opioid Therapy in the medical record. (PQRS 414)	Numerator	Patients evaluated for risk of misuse of opiates by using a brief validated instrument (e.g., Opioid Risk Tool, SOAAP-R) or patient interview at least once during opioid therapy	Having the risk evaluation added to the EMR workflow may be more difficult than the other measures.
				Denominator	All patients 18 and older prescribed opiates for longer than six weeks duration	The Opioid Rx will likely be in the EMR meds list; however, it may be hard to know if they have been taking them for 6 weeks